

CASE REPORT

Post-LASIK Ectasia associated with Pigmentary Glaucoma: Tomographic and Biomechanical Characterization

¹Thiago JMM Mazzeo, ²Nelson B Sena Jr, ³Ana LC Canedo, ⁴Isaac Ramos, ⁵Renata S da Silva
⁶Giovanni Colombini, ⁷Renato Ambrósio Jr

ABSTRACT

Aim: To report a case of bilateral post-laser *in situ* keratomileusis (LASIK) ectasia associated with pigmentary glaucoma (PG), in which intraocular pressure (IOP) assessment had been relentlessly underestimated by Goldmann's applanation tonometry (GAT).

Materials and methods: Case report and prospective literature review.

Results: The patient presented for second opinion regarding ectasia in the right eye, while ectasia was diagnosed in the left eye. Uncorrected visual acuity was 20/40 oculus dexter (OD) and 20/200 oculus sinister (OS). Distance corrected visual acuity was 20/20 in OD and 20/50 OS; GAT was 18/18 mm Hg. Slit-lamp biomicroscopy was relevant for a well-positioned superior hinge corneal flap and Krukenberg's spindle in both eyes, and Fleischer's ring in the left eye. Corneal compensated IOP was 47.8 and 43.8 mm Hg in OD and OS as measured by the ocular response analyzer (ORA; Reichert, Buffalo, New York, USA). The biomechanical compensated (IOPb) was 62.9 mm Hg OD and unmeasured OS by Corvis ST (Oculus; Wetzlar, Germany), which also demonstrated attenuated corneal deformation OU. Pentacam confirmed corneal ectasia in both eyes, being considerably worse in OS. One day after initiation of a fixed combination of beta blocker and carbonic anhydrase inhibitor, the patient noted significant improvement of vision with reduction of IOPb to 13 and 13.5 mm Hg in OD and OS respectively. Patient referred improvement in quality of vision and there was marked reduction on corneal scatter, thickness, and curvature.

Conclusion: Pigment dispersion syndrome (PDS) and PG may coexist with corneal ectasia. Careful consideration of the corneal impact on IOP assessment is mandatory when evaluation patients after laser vision correction (LVC). The IOP normalization may improve corneal ectasia in patients with coexisting glaucoma or ocular hypertension.

Keywords: Ectasia, Keratoconus, Pigmentary glaucoma.

How to cite this article: Mazzeo TJMM, Sena NB Jr, Canedo ALC, Ramos I, da Silva RS, Colombini G, Ambrósio R Jr. Post-LASIK Ectasia associated with Pigmentary Glaucoma: Tomographic and Biomechanical Characterization. *Int J Kerat Ect Cor Dis* 2018;7(1):61-65.

Source of support: Nil

Conflict of interest: None

INTRODUCTION

Corneal ectasia is an uncommon but serious vision-threatening complication of keratorefractive surgery.¹ It is characterized by a progressive topographic steepening associated with deterioration of visual acuity. As the number of patients who have refractive surgery increases, more cases of ectasia are being recognized.

Iatrogenic ectasia usually occurs within the first year of surgery, although cases have been reported up to 10 years postoperatively.² From the reported cases, a variety of risk factors were proposed, including younger age, high myopia, low residual stromal bed thickness, and low preoperative corneal thickness altering corneal biomechanics.³

Even though the role of corneal biomechanical properties in ectatic corneal diseases is well established, studies suggest that it also has a crucial role in glaucoma. This is because of the relation of corneal properties and the accuracy of IOP measurement, as well as the inherent risk of glaucomatous damage to the optic nerve.⁴ Corneal biomechanical factors can influence the accuracy of GAT and other tonometers. Although valuable in estimating glaucoma risk, central corneal thickness (CCT) is a suboptimal surrogate for the mechanical bending characteristics of the cornea, and it is but one component of a complex interaction between the viscous and elastic properties of the cornea, which ultimately combine to influence IOP measurements in not entirely predictable ways.^{5,6}

^{1,2}Resident, ³⁻⁵Research Associate, ⁶Professor and Chair
⁷Founder, Scientific Director, Adjunct Professor, and Affiliated Professor

^{1,2,6}Department of Ophthalmology, Federal University of the State of Rio de Janeiro, Rio de Janeiro, Brazil

³⁻⁵Rio de Janeiro Corneal Tomography and Biomechanics Study Group, Rio de Janeiro, Brazil

⁷Department of Ophthalmology, Federal University of the State of Rio de Janeiro, Rio de Janeiro, Brazil; Rio de Janeiro Corneal Tomography and Biomechanics Study Group, Rio de Janeiro Brazil, Department of Ophthalmology, Federal University of São Paulo, São Paulo, Brazil

Corresponding Author: Renato Ambrósio Jr, Founder, Scientific Director, Adjunct Professor, and Affiliated Professor Department of Ophthalmology, Federal University of the State of Rio de Janeiro, Rio de Janeiro, Brazil; de Janeiro Corneal Tomography and Biomechanics Study Group, Rio de Janeiro Brazil, Department of Ophthalmology, Federal University of São Paulo, São Paulo, Brazil, e-mail: dr.renatoambrosio@gmail.com

We report a case of bilateral corneal ectasia 9 years post-LASIK, coexisting with PG. The very high IOP had not been detected, as it was considerably underestimated by GAT. In addition, the case illustrates the impact of IOP on corneal transparency and geometry.

CASE REPORT

A 31-year-old man was referred due to progressive visual acuity loss for the last 6 months in OS due to post-LASIK ectasia and suspicious of ectasia in OD. The patient had no records of previous ocular disease and a negative family history for glaucoma or keratoconus.

The patient reported bilateral LASIK 9 years ago for treating moderate myopia. No detailed information was available regarding flap, ablation, or refractive treatment, nor preoperative corneal parameters.

Uncorrected visual acuity was 20/40 OD and 20/200 OS. Distance corrected visual acuity was 20/20 (-2.00 -1.50 x 160) in OD and 20/50 (1.25 -3.75 x 125) in OS.

At the slit-lamp biomicroscopy, patient had a well-positioned and clear LASIK flap with superior hinge in both eyes. There was very mild corneal edema and Krukenberg's spindles on both eyes (Fig. 1). Fleisher's ring was evident in OS; IOP (GAT) was 18 mm Hg in both eyes (2:15 pm). Corneal compensated IOP was 47.8 and 43.8 mm Hg in OD and OS respectively, as measured by the ORA (Reichert, Buffalo, New York, USA). The IOPb was 62.9 mm Hg OD and unmeasured OS by Corvis ST (Oculus; Wetzlar, Germany), which also demonstrated




Fig. 1: Right eye presenting mild corneal edema and Krukenberg's spindle. Margin of LASIK flap is also noticeable

attenuated corneal deformation OU (Fig. 2). Pentacam HR corneal tomography (Oculus; Wetzlar, Germany) showed corneal ectasia in both eyes, but considerably worse in OS (Figs 3A and 4A). Ectasia OD was confirmed based on the posterior elevation, with an elevation of 21 μm the thinnest point using the best fit sphere for 8 mm. Fundoscopic exam showed a linear cup disk/ratio of 0.3 OD and 0.7 OS.

The diagnosis of PG associated with corneal ectasia was established. The patient was asked to return 1 day after initiation of a fixed combination of beta blocker and carbonic anhydrase inhibitor [Azorga® (brinzolamide 1% and timolol maleate 0.5%); Alcon; Fort Worth, TX, US].






Fig. 2: Vinciguerra screening report from corvis ST measuring a 71.5 mm Hg IOP in OD, with a very low corneal amplitude deformation



Figs 3A to F: Pentacam OD. Curvature after (A), prior (B) to IOP reduction, and subtraction (C); and thickness after (D), prior (E) to IOP reduction, and subtraction (F)



Figs 4A to F: Pentacam OS. Curvature after (A), prior (B) to IOP reduction, and subtraction (C); and thickness after (D), prior (E) to IOP reduction, and subtraction (F)

The patient noted significant improvement of vision with uncorrected distance visual acuity of 20/25 OD and 20/80 OS. Manifest refraction was not performed at this visit. There was marked reduction of IOPb to 13 and 13.5 mm Hg in OD and OS respectively, with deformation corneal response consistent with the diagnosis of ectasia. Figure 5 demonstrates the Corvis ST data of the right eye in the second visit.

Marked reduction on corneal scatter, thickness, and curvature was noted in both eyes (Figs 3 and 4). Corneal appplanation was 2.2D in OD (Fig. 3C) and 5.4D in OS (Fig. 4C). Corneal thinning (reduction of edema) was noted in both eyes (Figs 3F and 4F).

The patient was advised to consider cross-linking in both eyes and keep medical treatment and clinical control for glaucoma. The patient returned to his original city and no follow-up was further obtained.

DISCUSSION

Iatrogenic corneal ectasia is a rare, but very severe complication of LASIK.⁷ There are recognized risk factors for the development of ectasia after keratorefractive surgery, including preoperative mild (or forme fruste) keratoconus, thin corneal pachymetry, thick LASIK flap, high corrections, and younger age.¹ But other factors are




Fig. 5: Vinciguerra screening report from corvis ST OD after IOP reduction

associated with ectasia progression, such as eye rubbing and pregnancy.^{1,8}

In fact, there are different mechanisms for ectasia to develop following keratorefractive surgery, including the inherent corneal predisposition to manifesting ectasia and the weakening by the impact from the procedure.⁹ The role of postoperative trauma, such as eye rubbing and high IOP may also be very relevant for the biomechanical decompensation.¹⁰

While biomechanical properties of the cornea are of paramount importance in the keratorefractive surgery evaluation, it also has been related to glaucoma field. This is because of its relation with IOP measurement, but also due to the relation to the risk of glaucomatous damage to the optic nerve. The ORA and Corvis ST are noncontact tonometers that use a pulse of air to apply force to applanate the cornea and measure the IOP and try calculating a corrected IOP value, considering corneal deformation properties.¹¹

Several studies investigating the potential utility of the corneal hysteresis measurement in glaucoma have been published, and it seems that low corneal hysteresis, but not CCT or IOP, is associated with progressive glaucomatous optic neuropathy.^{5,12}

The most common and accepted method for measuring IOP for over 50 years has been the GAT. Nevertheless, as we better understand corneal biomechanics, its imitations are we increasingly appreciated.¹³ Algorithms to correct for CCT to estimate IOP may oversimplify the effects of corneal biomechanics.

Corneal thickness is but one component of a complex interaction between the viscous and elastic properties of the cornea, which ultimately combine to influence IOP measurements in not entirely predictable ways. And even so, as a solo biomechanical component, it may not be reliable, since different corneal responses have been observed in corneas having similar thickness with the same deformation stimuli.^{5,11}

Many authors have found an artificial underestimation of IOP measurements by different types of tonometers in eyes that underwent LVC.^{14,15} As the number of patients who have LVC increases each day, falsely low IOP due to IOP measuring biomechanical limitations could precariously postpone the diagnosis of glaucoma or can interfere in the adequate control of the disease.¹⁶

The ophthalmologist who exam post-LASIK patients has to maintain a high index of suspicion for possible ocular hypertension and glaucoma patients, even if the IOP reading is low. The degree of myopic correction (which translates to ablation depth) may seem to be related to reduction in post-LASIK IOP. However, further studies are needed to determine if a critical level of myopic correction is required before a significant change in IOP post-LASIK occurs.¹⁶

Studies observed that patients with PDS, 5 to 10% will develop PG 5 to 6 years after diagnoses, 15% after 15 years, and 35% after 35 years. As the quiescent phase of PG occurs, the increased angle pigmentation tends to clear, and with this, the reversal of the pigmentary

signs in older patients can lead to a misdiagnosis of primary open-angle glaucoma or normal tension glaucoma when associated with IOP normalization. It also seems that the degree of myopia in PDS patients that develop PG is higher than the degree of myopia in those who do not develop.¹⁷ Interestingly, the relationship between PDS and corneal ectatic disease has not been established yet.

This article illustrates a case of ectasia that was associated with PG in which GAT limitations limited the ability for the diagnosis. This is true that a more careful slit lamp evaluation of the cornea and a comprehensive analysis including fundus exam would provide more information that would facilitate the diagnosis. Nevertheless, the role of novel methods for IOP assessment is unquestionably emphasized in this report. Refractive surgeons must be aware that there are bidirectional influences of IOP and corneal properties assessment.

REFERENCES

1. Binder PS. Analysis of ectasia after laser *in situ* keratomileusis: risk factors. *J Cataract Refract Surg* 2007 Sep;33(9):1530-1538.
2. Randleman JB, Trattler WB, Stulting RD. Validation of the ectasia risk score system for preoperative laser *in situ* keratomileusis screening. *Am J Ophthalmol* 2008 May;145(5):813-818.
3. Giri P, Azar DT. Risk profiles of ectasia after keratorefractive surgery. *Curr Opin Ophthalmol* 2017 Jul;28(4):337-342.
4. Broman AT, Congdon NG, Bandeen-Roche K, Quigley HA. Influence of corneal structure, corneal responsiveness, and other ocular parameters on tonometric measurement of intraocular pressure. *J Glaucoma* 2007 Oct-Nov;16(7):581-588.
5. Clement CI, Parker DG, Goldberg I. Intra-ocular pressure measurement in a patient with a thin, thick or abnormal cornea. *Open Ophthalmol J* 2016 Feb;10:35-43.
6. Duch S, Serra A, Castanera J, Abos R, Quintana M. Tonometry after laser *in situ* keratomileusis treatment. *J Glaucoma* 2001 Aug;10(4):261-265.
7. Haw WW, Manche EE. Iatrogenic keratectasia after a deep primary keratotomy during laser *in situ* keratomileusis. *Am J Ophthalmol* 2001 Dec;132(6):920-921.
8. Spadea L, Cantera E, Cortes M, Conocchia NE, Stewart CW. Corneal ectasia after myopic laser *in situ* keratomileusis: a long-term study. *Clin Ophthalmol* 2012 Nov;6:1801-1813.
9. Santhiago MR. Percent tissue altered and corneal ectasia. *Curr Opin Ophthalmol* 2016 Jul;27(4):311-315.
10. Deol M, Taylor DA, Radcliffe NM. Corneal hysteresis and its relevance to glaucoma. *Curr Opin Ophthalmol* 2015 Mar;26(2):96-102.
11. Ambrosio JR, Ramos I, Luz A, Faria FC, Steinmueller A, Krug M, Belin MW, Roberts CJ. Dynamic ultra high speed Scheimpflug imaging for assessing corneal biomechanical properties. *Braz J Ophthalmol* 2013 Mar-Apr;72(2):99-102.
12. Hirasawa K, Matsuura M, Murata H, Nakakura S, Nakao Y, Kiuchi Y, Asaoka R. Association between corneal biomechanical properties with ocular response analyzer and also CorvisST tonometry, and glaucomatous visual field severity. *Transl Vis Sci Technol* 2017 Jun;6(3):18.
13. Correia FF, Ramos I, Roberts CJ, Steinmueller A, Krug M, Ambrosio R Jr. Impact of chamber pressure and material properties on the deformation response of corneal models measured by dynamic ultra-high-speed Scheimpflug imaging. *Arq Bras Oftalmol* 2013 Oct;76(5):278-281.
14. Cronemberger S, Guimaraes CS, Calixto N, Calixto JM. Intraocular pressure and ocular rigidity after LASIK. *Arq Bras Oftalmol* 2009 Jul-Aug;72(4):439-443.
15. Razeghinejad MR, Nouri-Mahdavi K, Perera S. Primary open angle glaucoma and post-LASIK keratectasia. *J Ophthalmic Vis Res* 2010 Jul;5(3):196-201.
16. Tsai AS, Loon SC. Intraocular pressure assessment after laser *in situ* keratomileusis: a review. *Clin Exp Ophthalmol* 2012 Apr;40(3):295-304.
17. Niyadurupola N, Broadway DC. Pigment dispersion syndrome and pigmentary glaucoma—a major review. *Clin Exp Ophthalmol* 2008 Dec;36(9):868-882.